

Periodontal Associates

OF MEMPHIS

Periodontal, Laser and Dental Implant Therapy

6268 Poplar Avenue · Memphis, TN 38119 · phone 901-761-3770 · toll free 800-824-1628

fax 901-761-3775 www.PerioMem.com Info@PerioMem.com

Roger D. Craddock, D.D.S.
Mitchel S. Godat, D.D.S., M.S.
Board Certified Periodontist

Partners Emeritus:
Preston D. Miller, Jr., D.D.S.
James R. Ross, D.D.S., M.S.

POST- ESTHETIC CROWN LENGTHENING, SOFT TISSUE GRAFT, RIDGE AUGMENTATION, FRENECTOMY, WILCKODONTICS OR ORTHODONTIC UNCOVERY INSTRUCTIONS

Please read general Pre & Post Appointment Instructions. These are additional notes following the above procedures.

- 1) **If you had a bone or soft tissue graft** note the two major reasons for graft failure are:
- 2) Smoking during the healing phase
- 3) Inadvertently hitting the graft with a toothbrush before it has had a chance to attach to the root.

Following Esthetic crown lengthening you may notice spacing between your teeth after surgery. These spaces will fill in with gum tissue as you heal.

Please comply with the following:

- 1) Do not pull on your lips to look at the treated area. This could dislodge sutures and decrease the success of your treatment especially if you have had a hard or soft tissue graft.
- 2) Do not bite into any foods. This could damage the healing gum tissue. Please cut food into small pieces. (See food instructions under general instructions.)
- 3) After 5 days, you may gently clean your teeth with a Q-tip. Do not clean the gum area or you could cause the graft to fail. Clean the rest of your mouth as normal.
- 4) Do not exercise vigorously until you come back for your post-op appointment.
- 5) Refrain from kissing and playing a mouth musical instrument until your post-op appointment.
- 6) Do not wear a removable appliance (partial or denture) without having the doctor adjust or relin it. Doing so could cause your graft to fail.
- 7) If wearing a removable appliance, do not remove it for 24 hours. After 24 hours, remove the appliance and rinse gently. Replace the appliance immediately. Continue this daily until completion of treatment. This prevents swelling of the tissue. If tissue swelling occurs with the appliance out of the mouth, you may not be able to wear the appliance for several days. Do not use denture adhesive unless approved by your doctor.

If you have had a bone or soft tissue graft, your sutures will probably be removed at your 2nd follow up appointment.

(Last updated March 28, 2011)

Read, Sign, and Send all consent forms at least 3 to 5 days before surgery.

Fax: 1.901.761.3775



Active Member
American Academy of Periodontology
Specialist in Periodontics

